

Jean Daniel Morel¹, Rubens Manoel dos Santos¹, Marco Aurélio Leite Fontes¹, Paulo Oswaldo Garcia², Fernanda Maria de Souza¹

FLORISTIC COMPARISON BETWEEN TWO TREE COMMUNITIES ASSOCIATED WITH HABITAT DESCRIPTOR VARIABLES

Keywords:
Plant ecology
Geoecology
Floristic composition,
Environmental variables

Histórico:
Recebido 01/12/2014
Aceito 15/11/2015

ABSTRACT: The knowledge about the influence of habitat variables is essential to understand the underlying ecological patterns in vegetation. This study compared the floristic composition of two forest communities located in different altitudes. Associated with this comparison, we used a methodology where habitat descriptor variables were scaled and interpreted by the biotic set sampled. We constructed one matrix with scores given to physical, biotic, vegetation, and anthropogenic variables in the field and one matrix with the species sampled and performed multivariate analyses. We found that the floristic communities differ between the different altitudes and that the methodology used showed significant variables for the ecological characterization of the sampled habitat.

COMPARAÇÃO FLORÍSTICA ENTRE DUAS COMUNIDADES ARBÓREAS ASSOCIADA A VARIÁVEIS DESCRITORAS DO AMBIENTE

Palavras chave:
Ecologia vegetal
Geoecologia
Composição florística
Variáveis ambientais

Correspondência:
morel.jean@gmail.com

RESUMO: O conhecimento sobre a influência das variáveis ambientais é essencial para que se compreenda os padrões ecológicos atuantes sobre a vegetação. Este trabalho comparou a composição florística de duas comunidades florestais localizadas em diferentes altitudes. Associada a essa comparação, foi utilizada uma metodologia onde variáveis descritoras do ambiente foram escalonadas e interpretadas junto ao conjunto biótico amostrado. Através de uma matriz, com a pontuação dada em campo para variáveis físicas, bióticas, da estrutura da vegetação e de antropização, e de outra, contendo as espécies amostradas, análises multivariadas foram utilizadas. Constatou-se que as comunidades diferem floristicamente entre as diferentes altitudes e que a metodologia utilizada indicou variáveis importantes para a caracterização ecológica dos ambientes amostrados.

DOI:

10.1590/01047760201521041934

¹ Universidade Federal de Lavras - Lavras, Minas Gerais, Brasil

² Instituto Federal de Educação, Ciência e Tecnologia do Sul de Minas Gerais - Muzambinho, Minas Gerais, Brasil

INTRODUCTION

The distribution of plant species is determined by physicochemical and biological factors, and by a combination of these factors (CONTI; FURLAN, 2009; COX; MOORE, 2011). At the macroscale, climate is the major determinant of plant distribution, as it provides different conditions of temperature and precipitation (WHITTAKER, 1975; WALTER, 1985). Conversely, variations in topography and soils are the main determinants of vegetation at smaller scales (NEWTON, 2007; FERREIRA JÚNIOR et al., 2009). Moreover, interactions between species can also affect plant distribution, as the establishment and growth of several species can be inhibited or favored depending on the distribution of other plant species (KENT; COKER, 1992; SANTOS et al., 2012). Natural or anthropogenic disturbance events also cause changes in vegetation and may cause changes in the structure and dynamics of plant communities and populations (BEGON et al., 2007; GHAZOU; SHEIL, 2010).

The influence of environmental variables on plant species may result in different distribution patterns that vary over time and space (MEIRELES et al., 2008; SANTOS et al., 2012). Variation in species distributions requires that the combined effects of vegetation and environmental variables be evaluated together (TER BRAAK, 1987), taking into account that some variables are more responsive than others when a particular group of species is considered. Thus, it is often necessary to combine methods with floristic surveys and forest structure studies to elucidate ecological patterns and processes. Nevertheless, some of the information that characterizes the habitat remains unknown due to measuring difficulties or the high cost of quantification.

This study aimed to evaluate the floristic variation of two plant communities at different altitudes from a qualitative method for habitat characterization that uses a set of scaled variables. We expect that this method may serve as support information for the evaluated communities and provides additional parameters to help explain the ecological patterns of vegetation.

MATERIAL AND METHODS

We listed characteristics of native forest habitats based on a literature review, considering biotic and abiotic factors, arboreal physiognomy, and human disturbances. We selected 20 variables that were treated as habitat descriptor variables (HDV). These variables were grouped into the categories described below. A Abiotic descriptors: A1 Slope, A2 Rocky outcrops, A3 Leaf litter deposition, A4 Humidity, and A5 Wind exposure. B Biotic descriptors: B1 Bamboo density, B2 Bromeliad density, B3 Orchid density, B4 Bryophyte density, B5 Lichen density, and B6 Liana

density. C Arboreal/sub-arboreal vegetation descriptors: C1 Density of individuals, C2 Size of individuals, C3 Canopy stratification, C4 Canopy openness, and C5 Understory stratification. D Anthropization descriptors: D1 Abundance of cut stumps, D2 Abundance of trails, D3 Abundance of cattle feces, and D4 Abundance of fire evidence.

With the selected variables we constructed a semiquantitative matrix for survey in the field, as in the methods of phenological assessment (FOURNIER, 1974; WHEELWRIGHT, 1985). We used an interval scale (0 – 4) to infer the influence of variables on sampling units, with 0 indicating the lack of influence, 1 indicating a 1 – 25% influence, 2 to 26 – 50% influence, 3 to 51 – 75% influence, and 4 indicating a 76 – 100% influence.

Two areas were sampled in the same region, in the municipality of Itamonte, on a hillside of the Mantiqueira range, southern Minas Gerais state, Brazil. The floristic composition was sampled at 1500 m and 1900 m above sea-level (a.s.l.), in 30 sample points (15 in each area) delimited by a circle with 10 m radius, totaling 0,942 ha of sampling. At each sample point, all arboreal and sub-arboreal species were sampled, considering all individuals over 1 m high. Simultaneously to the vegetation survey, habitat descriptor variables were scaled using the habitat matrix.

Detrended Correspondence Analysis (DCA) (TER BRAAK, 1987) from a binary presence-absence species matrix and a categorical matrix of altitude was used to check the effect of altitude on the distribution of species independent of other habitat variables. Canonical Correspondence Analysis (CCA) (TER BRAAK, 1987) was used to infer the influence of HDVs on the distribution of species. In this case, in addition to the species matrix, we constructed a matrix with HDV data and used the Monte Carlo test to test the significance of correlations. CCA was performed on all habitat variables (CCA1) and on variables with correlation above 0.7 with the vegetation gradient (CCA2), considering the first two axes. The analyses were performed using PC-ORD 5.10 for Windows software (MCCUNE; MEFFORD, 2006).

RESULTS AND DISCUSSION

In total, 133 species were sampled, 98 at 1500 m and 78 at 1900 m. Of the total species sampled, 55 were unique to the lower altitude area and 35 were sampled in the higher altitude area only. These results are in accordance with the notion that species richness decreases with altitude in tropical forests (GENTRY, 1995; RAHBK, 1995; ADAMS, 2010; SCHEER et al., 2011). The list of species is shown in Table 1. In addition to species surveys, understanding the influence of habitat variables on the organization of biological communities helps corroborate the patterns observed and provides the delimitation of groups and the knowledge of the relationships between these groups and their habitats. The survey of the variables (Table 2) provides a greater basis for

Table 1 List of arboreal and sub-arboreal species in sampling points at two altitudes in the Mantiqueira range, municipality of Itamonte, Minas Gerais state, Brazil (0 = absence of the specie; 1 = presence of the specie).

Tabela 1 Lista de espécies arbóreas e subarbóreas verificadas nos pontos amostrais de duas cotas altitudinais da Serra da Mantiqueira no município de Itamonte, Minas Gerais, Brasil (0 = ausência da espécie; 1 = presença da espécie).

FAMILY / Specie	1900 m																
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15		
ANNONACEAE																	
<i>Annona cacans</i> Warm.	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0
<i>Annona sericea</i> Dunal	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0
<i>Cardiopetalum calophyllum</i> Schtdl.	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0
<i>Guatteria australis</i> A.St.-Hil.	1	0	1	1	1	0	1	1	1	1	0	1	0	0	1	0	0
<i>Guatteria sellowiana</i> Schtdl.	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
<i>Rollinia sylvatica</i> (A.St.-Hil.) Mart.	1	1	0	1	1	0	0	0	1	0	1	1	0	0	0	0	0
APOCYNACEAE																	
<i>Aspidosperma olivaceum</i> Müll.Arg.	1	1	1	1	0	0	0	1	1	1	0	0	0	0	0	0	0
AQUIFOLIACEAE																	
<i>Ilex conocarpa</i> Reissek	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Ilex paraguariensis</i> A.St.-Hil.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
ARALIACEAE																	
<i>Schefflera calva</i> (Cham.) Frodin & Fiaschi	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
ARAUCARIACEAE																	
<i>Araucaria angustifolia</i> (Bertol.) Kuntze	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ASTERACEAE																	
<i>Baccharis serrulata</i> DC.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
<i>Dasyphyllum tomentosum</i> (Spreng.) Cabrera	1	1	1	1	0	0	1	1	0	0	0	0	0	0	0	1	0
<i>Vernonanthura divaricata</i> (Spreng.) H.Rob.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0
BIGNONIACEAE																	
<i>Cybistax antisyphilitica</i> (Mart.) Mart.	1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0
<i>Handroanthus catarinensis</i> (A.H.Gentry) S.O.Grose	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0
BURSERACEAE																	
<i>Protium spruceanum</i> (Benth.) Engl.	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
CARDIOPTERIDACEAE																	
<i>Citronella paniculata</i> (Mart.) R.A.Howard	0	1	1	0	1	0	0	0	1	1	1	1	1	0	0	0	1

...be continue

...Continuation	1900 m														
	1500 m														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
FAMILY / Specie															
CELASTRACEAE															
<i>Maytenus evonymoides</i> Reissek	0	0	0	0	1	0	0	0	0	1	0	0	0	0	1
<i>Maytenus salicifolia</i> Reissek	0	1	1	0	0	0	0	1	0	1	0	0	0	0	1
CHLORANTHACEAE															
<i>Hedyosmum brasiliense</i> Miq.	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
CLETHRACEAE															
<i>Clethra scabra</i> Pers.	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0
CLUSIACEAE															
<i>Chrysochlamys saldanhae</i> (Engl.) Oliveira-Filho	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0
CUNONIACEAE															
<i>Lamanonia ternata</i> Vell.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CYATHEACEAE															
<i>Cyathea delgadii</i> Sternb.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ELAEOCARPACEAE															
<i>Sloanea monosperma</i> Vell.	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0
EUPHORBIACEAE															
<i>Alchornea triplinervia</i> (Spreng.) Müll.Arg.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Croton floribundus</i> Spreng.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Croton piptocalyx</i> Müll.Arg.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
FABACEAE															
<i>Bauhinia longifolia</i> (Bong.) D.Dietr.	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
<i>Coursekia rostrata</i> Benth.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Dalbergia brasiliensis</i> Vogel	0	1	1	0	0	1	1	0	1	0	0	0	0	0	0
<i>Inga platyptera</i> Benth.	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0
<i>Leucocylon incuriale</i> (Vell.) Barneby & J.W.Grimes	1	1	1	0	0	0	0	0	0	1	0	0	0	0	0
<i>Machaerium nycitans</i> (Vell.) Benth.	1	0	0	0	0	0	0	1	0	0	1	0	0	0	0
<i>Machaerium villosum</i> Vogel	0	0	0	0	0	0	0	0	1	1	1	0	0	0	0
<i>Ormosia fastigiata</i> Tul.	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0
<i>Platycamus regnellii</i> Benth.	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
<i>Senna splendida</i> (Vogel) H.S.Irwin & Barneby	0	1	1	0	0	1	0	0	0	1	0	0	0	0	0

...be continue

...Continuation

FAMILY / Specie	1500 m															1900 m																							
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15									
LACISTEMATACEAE																																							
<i>Lacistema hasslerianum</i> Chodat	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
LAURACEAE																																							
<i>Cryptocarya aschersoniani</i> Mez	0	0	0	1	0	1	0	1	1	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1
<i>Endlicheria paniculata</i> (Spreng.) J.F.Macbr.	1	1	0	1	1	1	1	1	1	0	1	1	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Nectandra nitidula</i> Nees	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Ocotea diospyrifolia</i> (Meisn.) Mez	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1
<i>Ocotea odorifera</i> (Vell.) Rohwer	0	0	0	1	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Ocotea pulchella</i> Mart.	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Persea major</i> L.E.Kopp	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Persea willdenowii</i> Kosterm.	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MAGNOLIACEAE																																							
<i>Magnolia ovata</i> (A.St.-Hil.) Spreng.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MALPIGIACEAE																																							
<i>Byrsonima laxiflora</i> Griseb.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MALVACEAE																																							
<i>Abutilon</i> sp.	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Luehea divaricata</i> Mart.	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Pseudobombax grandiflorum</i> (Cav.) A.Robyns	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Pseudobombax longiflorum</i> (Mart. & Zucc.) A.Robyns	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MELASTOMATACEAE																																							
<i>Leandra aurea</i> (Cham.) Cogn.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0	1	0	1
<i>Leandra quinquevata</i> (DC.) Cogn.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0	1	1
<i>Miconia castaneiflora</i> Naudin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	1	0	1	0	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0
<i>Miconia corallina</i> Spring	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Miconia latecrenata</i> (DC.) Naudin	1	1	1	1	1	1	1	1	1	1	0	0	1	1	0	1	1	0	0	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
<i>Miconia sellowiana</i> Naudin	1	1	1	1	1	1	1	1	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
<i>Miconia trianae</i> Cogn.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Tibouchina estrellensis</i> (Raddi) Cogn.	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MELIACEAE																																							

...be continue

...Continuation

FAMILY / Specie	1500 m															1900 m														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<i>Myrciaria floribunda</i> (H.West ex Willd.) O.Berg	1	1	0	1	0	1	0	1	0	1	0	1	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0
<i>Pimenta pseudocaryophyllus</i> (Gomes) Landrum	1	1	1	1	1	0	0	1	1	1	0	1	1	1	1	1	0	0	1	1	1	1	1	1	1	1	0	1	1	0
<i>Psidium cattleianum</i> Sabine	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0
<i>Psidium rufum</i> DC.	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0
<i>Siphoneugena densiflora</i> O.Berg	0	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Siphoneugena reitzii</i> D.Legrand	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Siphoneugena widgreniana</i> O.Berg	1	1	1	1	1	1	1	0	0	0	1	1	1	1	1	0	1	0	1	0	1	1	0	1	1	1	1	1	1	1
OCHNACEAE																														
<i>Ouratea floribunda</i> Engl.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	0	1	0	0	1	1	1
OPLIACEAE																														
<i>Agonandra excelsa</i> Griseb.	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
PERACEAE																														
<i>Pera glabrata</i> (Schott) Poepp. ex Baill.	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
PIPERACEAE																														
<i>Piper aduncum</i> L.	1	1	1	1	1	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Pothomorphe umbellata</i> L. Miq	0	0	1	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
PROTEACEAE																														
<i>Euplassa legalis</i> (Vell.) I.M.Johnst.	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Roupala montana</i> Aubl.	1	1	0	1	0	1	1	0	1	1	1	1	1	1	1	1	0	1	0	1	0	1	0	0	0	1	0	0	0	0
<i>Roupala rhombifolia</i> Mart. ex Meisn.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
ROSACEAE																														
<i>Prunus myrtifolia</i> (L.) Urb.	1	1	1	1	0	0	1	1	1	1	0	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0
RUBIACEAE																														
<i>Chomelia sericea</i> Müll.Arg.	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Cordia sessilis</i> (Vell.) Kuntze	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	1	1	1	1	1	1	1	1	1	1	1	1	0
<i>Guettarda uruguensis</i> Cham. & Schtdl.	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Psychotria deflexa</i> DC.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Psychotria hastisepala</i> Müll.Arg.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	1	0	0	1	0
<i>Psychotria stachyoides</i> Benth.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Psychotria suterella</i> Müll.Arg.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Psychotria vellosiana</i> Benth.	1	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

...be continue

...Continuation	1500 m															1900 m														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
FAMILY / Specie	0	1	0	1	1	1	1	0	1	1	1	1	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0
<i>Rudgea jasminoides</i> (Cham.) Müll.Arg.																														
SALICACEAE																														
<i>Casearia decandra</i> Jacq.	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Casearia lasiophylla</i> Eichler	1	1	1	0	0	0	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Casearia obliqua</i> Spreng.	0	1	1	1	0	1	0	0	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Casearia sylvestris</i> Sw.	1	1	1	0	1	1	0	0	0	0	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Xylosma prockia</i> (Turcz.) Turcz.	1	1	1	1	1	1	1	0	0	1	1	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SAPINDACEAE																														
<i>Cupania vernalis</i> Cambess.	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Cupania zanthoxyloides</i> Cambess.	1	1	1	1	1	0	0	1	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SIPARUNACEAE																														
<i>Siparuna cujabana</i> (Mart.) A. DC.	0	0	0	1	1	0	0	0	0	1	0	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SOLANACEAE																														
<i>Brunfelsia uniflora</i> (Pohl) D. Don	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	1	1	0	1	1	0	0	0	0	0
<i>Cestrum schlechtendalii</i> G. Don	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0
<i>Solanum asperum</i> Rich.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Solanum leucodendron</i> Sendtn.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	1	0	1	0	1	0	0	1	0
<i>Solanum pseudoquina</i> A. St.-Hil.	1	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	1	1	0	0	0
SYMPLOCACEAE																														
<i>Symplocos affinis</i> Hort. ex Pasq.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Symplocos celastrina</i> Mart. ex Miq.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	1	0	0	0	0	1	0	1	1	1	1
<i>Symplocos insignis</i> Brand	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	1	0	1	1	1	1	1
THYMELAEACEAE																														
<i>Daphnopsis brasiliensis</i> Mart. & Zucc.	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Daphnopsis coriacea</i> Taub.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	0	1	1	0	1	0	1	0	0	1	0	0
VOCHYSIACEAE																														
<i>Vochysia magnifica</i> Warm.	0	1	1	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Vochysia tucanorum</i> Mart.	1	1	1	1	1	1	1	1	1	1	1	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
WINTERACEAE																														
<i>Drimys brasiliensis</i> Miels	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table 2 Habitat descriptor variables evaluated in points at two altitudes in the Mantiqueira range, municipality of Itamonte, Minas Gerais state, Brazil (0 = lack of influence; 1 = 1 – 25% influence; 2 = 26 – 50% influence; 3 = 51 – 75% influence; 4 = 76 – 100% influence).

Tabela 2 Variáveis descritoras do ambiente avaliadas nos pontos amostrais de duas cotas altitudinais da Serra da Mantiqueira no município de Itamonte, Minas Gerais, Brasil (0 = ausência de influência; 1 = 1 – 25% de influência; 2 = 26 – 50% de influência; 3 = 51 – 75% de influência; 4 = 76 – 100% de influência).

A Abiotic descriptors																															
A1 Slope																															
Altitude	1900 m																														
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Score	1	1	1	2	2	2	4	3	2	3	1	2	2	1	1	1	1	1	1	1	1	1	1	2	2	2	2	1	2	2	2
A2 Rocky outcrops																															
Altitude	1900m																														
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Score	1	0	1	0	1	1	1	0	1	2	1	3	3	2	0	0	3	0	0	0	0	0	0	0	0	0	2	1	0	0	
A3 Leaf litter deposition																															
Altitude	1900m																														
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Score	3	3	3	3	3	3	3	3	2	2	2	2	4	2	2	3	2	3	3	3	3	3	3	3	3	2	4	3	3	3	
A4 Humidity																															
Altitude	1500m																														
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Score	2	2	2	2	2	1	2	1	2	1	2	2	2	2	4	1	1	1	1	1	1	1	1	1	1	2	2	3	2	2	
A5 Wind exposure																															
Altitude	1500m																														
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Score	1	1	1	1	1	2	2	1	2	1	2	2	2	2	2	2	4	2	2	4	4	3	3	4	2	4	4	2	2	3	
B Biotic descriptors																															
B1 Bamboo density																															
Altitude	1900m																														
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
Score	0	0	0	0	0	3	4	4	4	4	1	0	0	1	1	0	0	4	3	3	3	4	3	3	4	2	2	3	2	0	0

...be continue

...Continuation

B2 Bromeliad density																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	0	0	0	0	0	0	0	0	0	0	1	1	2	1	1	1	1	1	1									
B3 Orchid density																												
A Abiotic descriptors																												
A1 Slope																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	0	1	1	1							
B4 Bryophyte density																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	1	2	2	2	2	1	2	2	1	2	2	1	4	3	3	2	2	3	3	1	3	3	1	3	4	4		
B5 Lichen density																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	1	2	1	1	1	1	1	2	2	2	2	2	2	3	2	4	3	4	3	3	3	3	3	3	3	3	4	
B6 Liana density																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	3	2	3	3	3	3	4	4	4	2	2	2	2	3	2	4	3	4	3	3	3	3	3	3	3	3	4	
C Arboreal/sub-arboreal vegetation descriptors																												
C1 Density of individuals																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	3	3	3	4	3	2	2	2	2	3	3	3	3	3	3	3	3	1	3	3	1	2	1	1	3	3	3	
C2 Size of individuals																												
1500m							1900m																					
Altitude																												
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15													
Score	2	3	3	3	3	2	2	2	2	2	3	3	3	3	3	3	2	2	2	2	2	2	2	2	2	2	2	3

...be continue

...Continuation

C3 Canopy stratification															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	3	3	3	3	2	2	2	3	3	3	3	3	2	2	3
C4 Canopy openness															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	1	1	2	2	2	2	2	2	2	2	3	4	2	2	2
C5 Understory stratification															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	4	3	3	3	2	1	1	1	2	3	3	3	3	3	4
D Anthropization descriptors															
D1 Abundance of cut stumps															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
D2 Abundance of trails															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
D3 Abundance of cattle feces															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
D4 Abundance of fire evidence															
1500m								1900m							
Altitude															
Point	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Score	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

inferences and better understanding of the influence of the variables used on vegetation patterns.

Ordination analysis showed that arboreal and sub-arboreal vegetation differed between altitudes (Figure 1). Axis 1, which represents the main floristic gradient, indicates that each altitude has a different group of species. The eigenvalue of this axis (0.558) shows a long gradient, indicating the occurrence of species unique to each altitude (TER BRAAK, 1995). Moreover, there is a greater cohesion in the group of species from 1900 m a.s.l. due to the presence of species typical of higher elevations such as *Byrsonima laxiflora*, *Drimys brasiliensis*, and *Symplocos celastrinea*, in addition to the genus *Ilex* (OLIVEIRA FILHO; FONTES, 2000).




Figure 1 Detrended Correspondence Analysis (DCA) diagram of the distribution of arboreal and sub-arboreal species sampled at two altitudes in the Mantiqueira range, municipality of Itamonte, Minas Gerais state, Brazil. A = points at 1500 m; and B = points at 1900 m.

Figura 1 Diagrama gerado pela Análise de Correspondência Retificada para a distribuição das espécies arbóreas e subarbóreas amostradas em duas cotas altitudinais da Serra da Mantiqueira no município de Itamonte, Minas Gerais, Brasil.

The difference in species composition between the two areas sampled was due to the differential distribution of species along the altitudinal gradient. Nevertheless, altitudinal variations comprise a complex set of variables related to temperature, air humidity, water availability, wind exposure, in addition to soil depth and drainage classes (CARVALHO et al., 2005; HOMEIER et al., 2010; BLUNDO et al., 2012).

Canonical Correspondence Analyses showed significant correlations between habitat variables and community organization (Table 3). All habitat descriptor variables (HDV), except for the four anthropogenic

variables (scored value 0), were used in CCA1, totaling 16 quantitative variables and one categorical variable (altitude). This analysis showed that the distribution of species correlated significantly with HDVs ($p < 0.01$).

Table 3 Summary of Canonical Correspondence Analysis (CCA) results of arboreal and sub-arboreal species sampled at two altitudes in the Mantiqueira range, municipality of Itamonte, Minas Gerais state, Brazil.

Tabela 3 Resumo dos resultados da Análise de Correspondência Canônica para as espécies arbóreas e subarbóreas amostradas em duas cotas altitudinais da Serra da Mantiqueira no município de Itamonte, Minas Gerais, Brasil.

	p	Axis 1	Axis 2	Axis 3
CCA1				
Monte Carlo test	0.001			
Eigenvalue		0.548	0.258	0.221
Species–environment correlation	0.996	0.976	0.978	
Cumulative percentage variance of species presence data	13.40	19.70	25.10	
Cumulative percentage variance of species–environment	18.80	27.20	34.90	
CCA2				
Monte Carlo test	0.001			
Eigenvalue		0.523	0.234	0.171
Species–environment correlation	0.978	0.958	0.969	
Cumulative percentage variance of species presence data	12.80	18.50	22.70	
Cumulative percentage variance of species–environment	42.80	61.10	75.60	

CCA2 was calculated using the five variables that correlated with the vegetation gradient ($r > 0.7$, considering the first two axes): humidity, wind exposure, bromeliad density, orchid density, and lichen density (Table 4). According to CCA2, axis 2 showed the correlation between humidity with the distribution of species sampled at 1500 m (Figure 2).

It should be noted that humidity, which is important for temperature regulation and nutrient transport in soil (CONTI; FURLAN, 2009), favors the establishment of species adapted to the microclimate provided by it. This effect was observed in the area at 1500 m, which was close to a water course. In fact, this variable probably favored the establishment of species such as *Casearia obliqua*, *Endlicheria paniculata*, *Luehea divaricata*, *Magnolia ovata*, *Platycomus regnellii*, *Protium spruceanum*, and *Vochysia magnifica*, whose

occurrences were previously associated with humid habitats (OLIVEIRA FILHO; FLUMINHAN FILHO, 1999; BOTREL et al., 2002; SOUZA et al., 2003; CARVALHO et al., 2005; ROCHA et al., 2005).

The variable wind exposure, most associated with the axis 1, is correlated to the distribution of species at 1900 m. This variable is associated with high altitudes and acts on the composition, structure and distribution of tree species in communities located in elevated areas (NEWTON, 2007; HOMEIER et al., 2010; BERTONCELLO et al., 2011). In study along a 2.5 km transect ranging in tropical montane cloud forests in north-western Costa Rica, Häger (2010) considered the wind exposure as a key variable to tree species turnover along environmental gradients on a local scale, therefore the constancy of the wind acts upon the air temperature and the cloudiness. Considering the ecology of cloud forests, the author also points out that these variables determine the occurrence of species, including endemic species, in habitat with strict conditions. (HÄGER, 2010).

Between the two communities evaluated the wind exposure should be acting in this way, contributing to the differentiation of the area located 1900 m a.s.l. In addition to the relationship with altitude wind exposure may also be associated with the occurrence of bamboos and canopy openness. Despite not entering in the inclusion criteria of CCA2, these two variables showed significant correlations with axis 1 to the area located at 1900 m.

Another variables more strongly associated with species distributions at 1900 m with greater amplitude along axis 2 were orchid, bromeliad and lichen density, all of which typical of montane forests (ADAMS, 2010; GHAZOUL; SHEIL, 2010; SCHEER et al., 2011). The presence of these botanicals groups in montane forests is related to the constant water supply provided by the cloudness, which makes its typical species of this habitat type (ADAMS, 2010; ACHARYA, 2011; BLUM et al., 2011; KRÖMER et al., 2013; DYMYTROVA et al., 2014). Thus, some tree species characteristics of high altitude as *Ilex paraguariensis*, *Croton piptocalyx*, *Leandra aurea*, *Ouratea floribunda*, *Roupala rhombifolia* e *Symplocos insignis*, exclusive from community located at 1900 m and sampled in majority points, may have their local occurrence related to microclimate conditions.

The use of HDVs explained a large part of the variance of the data. The cumulative variance of the first three axes in CCA2 was 22.7% for species data and 75.6% for the relationship between species and habitat variables (Table 3). Prado et al. (2002) argued that, in many cases, the first two or three axes explain a large

Table 4 Correlations between habitat variables and the first three axis in the Canonical Correspondence Analyses (CCA1) for communities located at two altitudes in the Mantiqueira range, municipality of Itamonte, Minas Gerais state, Brazil.

Tabela 4 Correlações entre variáveis ambientais e os três primeiros eixos da Análise de Correspondência Canônica (CCA1) para comunidades localizadas em duas altitudes na Serra da Mantiqueira, no município de Itamonte, Minas Gerais, Brasil.

Variable	Correlations		
	Axis 1	Axis 2	Axis 3
A1 Slope	0.256	-0.288	-0.056
A2 Rocky outcrops	0.390	-0.105	0.296
A3 Leaf litter deposition	0.040	-0.196	-0.295
A4 Humidity	0.301	0.783	0.155
A5 Wind exposure	-0.725	0.223	-0.014
B1 Bamboo density	-0.387	-0.090	-0.565
B2 Bromeliad density	-0.855	-0.092	0.015
B3 Orchid density	-0.857	-0.238	0.067
B4 Bryophyte density	-0.678	-0.385	-0.054
B5 Lichen density	-0.886	-0.058	0.219
B6 Liana density	0.249	-0.016	-0.467
C1 Density of individuals	0.358	-0.086	0.404
C2 Size of individuals	0.222	0.161	0.446
C3 Canopy stratification	0.164	-0.050	0.245
C4 Canopy openness	-0.612	0.103	-0.151
C5 Understory stratification	-0.299	0.026	0.483




Figure 2 Canonical Correspondence Analysis (CCA) diagram of the distribution of arboreal and sub-arboreal species sampled at two altitudes in the Mantiqueira range, municipality of Itamonte, Minas Gerais state, Brazil. A = points at 1500 m; and B = points at 1900 m.

Figura 2 Diagrama gerado pela Análise de Correspondência Canônica para a distribuição das espécies arbóreas e subarbóreas amostradas em duas cotas altitudinais da Serra da Mantiqueira no município de Itamonte, Minas Gerais, Brasil.

part of the variance (60–90%), which enables to use the results to describe the system without significant loss of information. Thus, even when the first two axes only are considered, the percentage values obtained in this study are high.

We consider that the use of HDVs provided parameters that allowed us to interpret and discuss the floristic composition of the sampled communities more widely, since in addition to the altitude, other habitat features were added. With the largest number of variables were obtained parameters allowed to extend the discussion. In addition to corroborate the influence of altitude the variables used showed characteristics of communities that extended their ecological interpretation.

ACKNOWLEDGMENTS

We are grateful to the owners and coordinators at RPPN Altomontana, where field work was conducted, and to students for their help with field work. We are also grateful to Coordination for the Improvement of Higher Education Personnel – CAPES, for the doctoral scholarship given to the first author, to Foundation for Research of Minas Gerais State – FAPEMIG, and the Post Graduate Program in Forestry at UFLA.

REFERENCES

- ACHARYA, K. P.; VETAAS, O. R.; BIRKS, H. J. B. Orchid species richness along Himalayan elevational gradients. **Journal of Biogeography**, Oxford, v. 38, n. 9, p. 1821-1833, set. 2011.
- ADAMS, J. **Vegetation-climate interaction: how plants make the global environment**. 2.ed. Chichester, UK: Springer Praxis Books, 2010. 266 p.
- BEGON, M.; TOWNSEND, C. R.; HARPER, J. L. **Ecologia: de indivíduos a ecossistemas**. 4. ed. Porto Alegre: Artmed, 2007. 740 p.
- BERTONCELLO, R.; YAMAMOTO, K.; MEIRELES, L. D.; SHEPHERD, G. J. A phytogeographic analysis of cloud forests and other forest subtypes amidst the Atlantic forests in south and southeast Brazil. **Biodiversity and Conservation**, Amsterdam, v. 20, n. 14, p. 3413-3433, dez. 2011.
- BLUM, C. T.; RODERJAN, C. V.; GALVÃO, F. Composição florística e distribuição altitudinal de epífitas vasculares da Floresta Ombrófila Densa na Serra da Prata, Morretes, Paraná, Brasil. **Biota Neotropica**, São Paulo, v. 11, n. 4, p. 141-149, out. 2011.
- BLUNDO, C.; MALIZIA, L. R.; BLAKE, J. G.; BROWN, A. D. Tree species distribution in Andean forests: influence of regional and local factors. **Journal of Tropical Ecology**, Cambridge, v. 28, n. 1, p. 83-95, jan. 2012.
- BOTREL, R. T.; OLIVEIRA FILHO, A. T. de; RODRIGUES, L.; CURI, N. Influência do solo e topografia sobre as variações da composição florística e estrutural da comunidade arbóreo-arbustiva de uma floresta estacional semidecidual em Ingá, MG. **Revista Brasileira de Botânica**, São Paulo, v. 25, n. 2, p. 195-213, 2002.
- CARVALHO, D. A.; OLIVEIRA FILHO, A. T.; BERG, E. V. D.; FONTES, M. A. L.; VILELA, E. A.; MARQUES, J. J. S. M.; CARVALHO, W. A. C. Variações florísticas e estruturais do componente arbóreo de uma floresta ombrófila altomontana às margens do rio Grande, Bocaina de Minas, MG, Brasil. **Acta Botanica Brasilica**, São Paulo, v. 19, n. 1, p. 91-109, jan./mar. 2005.
- CONTI, J. B.; FURLAN, S. A. Geoeecologia: o clima, os solos e a biota. In: ROSS, J. L. S. (Org.). **Geografia do Brasil**. 6. ed. São Paulo: Editora da Universidade de São Paulo, 2009. p. 67-208.
- COX, C. B.; MOORE, P. D. **Biogeografia: uma abordagem ecológica e evolucionária**. 7. ed. Rio de Janeiro: LTC, 2011. 398 p.
- DYMYTROVA, L.; NADYEINA, O.; HOBI, M. L.; CHEIDEGGER, C. Topographic and forest-stand variables determining epiphytic lichen diversity in the primeval beech forest in the Ukrainian Carpathians. **Biodiversity and Conservation**, Amsterdam, v. 23, n. 6, p. 1367-1394, jun. 2014.
- FERREIRA JÚNIOR, W. G.; SCHAEFER, C. E. G. R.; SILVA, A. F. Uma visão pedogeomorfológica sobre as formações florestais da Mata Altântica. In: MARTINS, S. V. (Ed.). **Ecologia de florestas tropicais do Brasil**. Viçosa: Editora UFV, 2009. cap. 4, p. 109-142.
- FOURNIER, L. A. Un metodo cuantitativo para la medición de características fenológicas en arboles. **Turrialba**, San José, v. 24, n. 4, p. 422-423, Oct./Dec. 1974.
- GHAZOUL, J.; SHEIL, D. **Tropical rain forest: ecology, diversity, and conservation**. Oxford: Oxford University Press, 2010. 516 p.
- GENTRY, A. H. Patterns of diversity and floristic composition in neotropical montane forest. In: CHURCHILL, S. P.; BASLEV, H.; FORERO, E.; LUTEYN, J. L. (Ed.). **Biodiversity and conservation of neotropical montane forest**. New York: The New York Botanical Garden Press, 1995. p. 103-126.
- HÄGER, A. The effect of climate and soil conditions on tree species turnover in a tropical montane cloud forest in Costa Rica. **Revista de Biología Tropical**, San José, v. 58, n. 4, p. 1489-1506, dez. 2010.

- HOMEIER, J.; BRECKLE, S. W.; GÜNTER, S.; ROLLENBECK, R. T.; LEUSCHNER, C. Tree diversity, forest structure and productivity along altitudinal and topographical gradients in a species-rich Ecuadorian Montane Rain Forest. **Biotropica**, Washington, v. 42, n. 2, p. 140-148, mar. 2010.
- KENT, M.; COKER, P. **Vegetation Description and Analysis: a practical approach**. London: Belhaven Press, 1992. 363 p.
- KRÖMER, T.; ACEBEY, A.; KLUGE, J.; KESSLER, M. Effects of altitude and climate in determining elevational plant species richness patterns: a case study from Los Tuxtlas, Mexico. **Flora**, Haale, v. 208, n. 3, p. 197-210, mar. 2013.
- MCCUNE, B.; MEFFORD, M. J. **PC-ORD: multivariate analysis of ecological data: version 5.1**. Gleneden Beach: MjM Software Design, 2006. 1 CD ROM.
- MEIRELES, L. D.; SHEPHERD, G. J.; KINOSHITA, L. S. Variações na composição florística e na estrutura fitossociológica de uma floresta ombrófila densa altomontana na Serra da Mantiqueira, Monte Verde, MG. **Revista Brasileira de Botânica, São Paulo**, v. 31, n. 4, p. 559-574, out./dez. 2008.
- NEWTON, A. C. **Forest Ecology and Conservation**. Oxford: Oxford University Press, 2007. 454 p.
- OLIVEIRA FILHO, A. T.; FLUMINHAN FILHO, M. Ecologia da vegetação do Parque Florestal Quedas do Rio Bonito. **Cerne**, Lavras, v. 5, n. 2, p. 51-64, 1999.
- OLIVEIRA FILHO, A. T.; FONTES, M. A. L. Patterns of floristic differentiation among Atlantic Forests in Southeastern Brazil and the influence of climate. **Biotropica**, Washington, v. 32, n. 4b, p. 793-810, dez. 2000.
- PRADO, P. I.; LEWINSOHN, T. M.; CARMO, R. L.; HOGAN, D. J. Ordenação multivariada na ecologia e seu uso em ciências ambientais. **Ambiente & Sociedade**, Campinas, v. 5, n. 10, p. 1-17, jan./jun. 2002.
- RAHBEK C. The elevational gradient of species richness: a uniform pattern? **Ecography**, Copenhagen, v. 18, n. 4, p. 200-205, dez. 1995.
- ROCHA, C. T. V.; CARVALHO, D. A.; FONTES, M. A. L.; OLIVEIRA FILHO, A. T.; BERG, E. V. D.; MARQUES, J. J. G. S. M. Comunidade arbórea de um continuum entre floresta paludosa e de encosta em Coqueiral, Minas Gerais, Brasil. **Revista Brasileira de Botânica**, São Paulo, v. 28, n. 2, p. 203-218, 2005.
- SANTOS, S. C.; BUDKE, J. C.; MULLER, A. Regeneração de espécies arbóreas sob a influência de *Merostachys multiramea* Hack. (Poaceae) em uma floresta subtropical. **Acta Botanica Brasilica**, Feira de Santana, v. 26, n. 1, p. 218-229, jan./mar. 2012.
- SCHERER, M. B.; MOCOCHINSKI, A. Y.; RODERJAN, C. V. Estrutura arbórea da Floresta Ombrófila Densa Altomontana de serras do Sul do Brasil. **Acta Botanica Brasilica**, Feira de Santana, v. 25, n. 4, p. 735-750, out./dez. 2011.
- SOUZA, J. dos S.; ESPÍRITO-SANTO, F. del bon; FONTES, M. A. L.; OLIVEIRA FILHO, A. T. de; BOTEZELLI, L. Análise das variações florísticas e estruturais da comunidade arbórea de um fragmento de floresta semidecídua às margens do rio Capivari, Lavras-MG. **Revista Árvore**, Viçosa, v. 27, n. 2, p. 185-206, 2003.
- TER BRAAK, C. J. F. The analysis of vegetation environment relationships by canonical correspondence analysis. **Vegetatio**, The Hage, v. 69, n. 3, p. 69-77, 1987.
- TER BRAAK, C. J. F. Ordination. In: JONGMAN, R. H. G.; TER BRAAK, C. J. F.; TONGEREN, O. F. R. van (Ed.) **Data analysis in community and landscape ecology**. Cambridge: University Press Cambridge, 1995. p. 91-173.
- WALTER H. **Vegetation of the earth and ecological systems of the geo-biosphere**. 3. ed. Berlin: Springer-Verlag, 1985. 318 p.
- WHEELWRIGHT, N. T. Competition for dispersers, and the timing of flowering and fruiting in a guild of tropical trees. **Oikos**, Copenhagen, v. 44, n. 3, p. 465-477, Dec. 1985.
- WHITTAKER, R. H. **Communities and Ecosystems**, 2. ed. New York: Macmillan, 1975. 385 p.

